



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Grasshopper Sparrow.	Cerulean Warbler.
White-crowned Sparrow.	Bay-breasted Warbler.
White-throated Sparrow.	Blackburnian Warbler.
Chipping Sparrow.	Black-throated Green Warbler.
Field Sparrow.	Palm Warbler.
Song Sparrow.	Oven-bird.
Swamp Sparrow.	Water-Thrush.
Towhee.	Louisiana Water-Thrush.
Cardinal.	Maryland Yellow-throat.
Rose-breasted Grosbeak.	Yellow-breasted Chat.
Indigo Bunting.	Wilson Warbler.
Scarlet Tanager.	Canadian Warbler.
Purple Martin.	American Redstart.
Cliff Swallow.	American Pipit.
Barn Swallow.	Catbird.
Tree Swallow.	Brown Thrasher.
Rough-winged Swallow.	House Wren.
Bank Swallow.	Winter Wren.
Migrant Shrike.	Long-billed Marsh Wren.
Red-eyed Vireo.	Brown Creeper.
Warbling Vireo.	White-breasted Nuthatch.
Yellow-throated Vireo.	Red-breasted Nuthatch.
Blue-headed Vireo.	Tufted Titmouse.
Nashville Warbler.	Chickadee.
Golden-winged Warbler.	Ruby-crowned Kinglet.
Black and white Warbler.	Blue-gray Gnatcatcher.
Blue-winged Warbler.	Wood Thrush.
Tennessee Warbler.	Wilson Thrush.
Yellow Warbler.	Olive-backed Thrush.
Black-throated Blue Warbler.	Gray-cheeked Thrush.
Myrtle Warbler.	Robin.
Magnolia Warbler.	Bluebird.
Chestnut-sided Warbler.	

WINTER BIRDS.

BY LYND S JONES.

The latitudinal variation in the range of many species of birds in winter is one of the problems which nothing but careful and persistent field work will solve. Sufficient interest attaches to the study to make the effort of determining what birds are about us in winter well worth the while. Bird studies in winter are not so hedged about with difficulties and inconveniences as study in August, because then dense foliage, excessive heat, insects and the timidity of the birds combine to thwart the purposes of the bird student. In winter there is only the cold and snow, which should not be serious hindrances to a healthy body.

WINTER BIRDS OF SCIOTO AND PIKE COUNTIES, OHIO.

For the sake of comparison with a study of the winter birds in Lorain county, this list is given for the extreme southern part of Ohio, almost directly south of Lorain county. The comparison brings out many interesting questions. It is well known that the southern part of the state experiences lower temperatures than the northern, during some part of the winter, but it also has less snow and a shorter time during which the food of the birds is covered. This would be reason enough why many of the birds which are not found in the northern counties during the winter find the southern counties fairly comfortable winter quarters. Rev. W. F. Henninger contributes this list:

SPECIES WHICH ARE RESIDENT, 42.

- Pied-billed Grebe, *Podilymbus podiceps*. Fairly common.
 Mallard, *Anas boschas*. Common.
 Killdeer, *Ægialitis vocifera*. Very common.
 Bob-white, *Colinus virginianus*. Very common.
 Ruffed Grouse, *Bonasa umbellus*. Common.
 Mourning Dove, *Zenaidura macroura*. Very common.
 Turkey Vulture, *Cathartes aura*. Common.
 Cooper Hawk, *Accipiter cooperi*. Common.
 Sharp-shinned Hawk, *Accipiter velox*. Not common.
 Red-tailed Hawk, *Buteo borealis*. Common.
 Red-shouldered Hawk, *Buteo lineatus*. Not common.
 Am. Sparrow Hawk, *Falco sparverius*. Very common.
 Barn Owl, *Strix pratincola*. Common.
 Long-eared Owl, *Asio wilsonianus*. Rare.
 Barred Owl, *Syrnium nebulosum*. Common.
 Screech Owl, *Megascops asio*. Very common.
 Great Horned Owl, *Bubo virginianus*. Common.
 Belted Kingfisher, *Ceryle alcyon*. Fairly common.
 Hairy Woodpecker, *Dryobates villosus*. Fairly common.
 Downy Woodpecker, *Drobates pubescens medianus*. Common.
 Northern Pileated Woodpecker, *Ceophlœus pileatus abieticola*.
 Rather rare.
 Red-bellied Woodpecker, *Melanerpes carolinus*. Common.
 Northern Flicker, *Colaptes auratus luteus*. Common.
 Prairie Horned Lark, *Otocorys alpestris praticola*. Common.
 Blue Jay, *Cyanocitta cristata*. Not common.
 American Crow, *Corvus americanus*. Common.
 Meadowlark, *Sturnella magna*. Common.
 American Goldfinch, *Astragalinus tristis*. Common.
 Chipping Sparrow, *Spizella socialis* (at times). Common.
 Song Sparrow, *Melospiza melodia*. Common.
 Towhee, *Pipilo erythrophthalmus*. Males only. Common.
 Cardinal, *Cardinalis cardinalis*. Very common.
 Cedar Waxwing, *Ampelis cedrorum*. Common.
 Migrant Shrike, *Lanius ludovicianus migrans*. Fairly common.

Carolina Wren, *Thryothorus ludovicianus*. Common.
 Bewick Wren, *Thryomanes bewickii*. Common.
 Chickadee, *Parus atricapillus*. Common.
 White-breasted Nuthatch, *Sitta carolinensis*. Common.
 Tufted Titmouse, *Parus bicolor*. Common.
 Carolina Chickadee, *Parus carolinensis*. Not common.
 American Robin, *Merula migratoria*. Common.
 Bluebird, *Sialia sialis*. Common.

WINTER RESIDENTS ONLY, 10.

Marsh Hawk, *Circus hudsonius*. Common.
 Am. Rough-legged Hawk, *Archibuteo lagopus sancti-johannis*.
 Rather rare.
 Bald Eagle, *Haliaeetus leucocephalus*. Rather rare.
 Osprey, *Pandion haliaetus carolinensis*. Fairly common.
 Short-eared Owl, *Asio accipitrinus*. Common.
 Rusty Blackbird, *Scolecophagus carolinus*. Common.
 White-throated Sparrow, *Zonotrichia albicollis*. Very common.
 Tree Sparrow, *Spizella monticola*. Very common.
 Slate-colored Junco, *Junco hyemalis*. Very comon.
 Northern Shrike, *Lanius borealis*. Rare.

RARE TRANSIENTS SOMETIMES SEEN IN JANUARY AND
 FEBRUARY, 8.

Herring Gull, *Larus argentatus*. Rare.
 Red-breasted Merganser, *Merganser serrator*. One record only.
 Golden Eagle, *Aquila chrysaetos*. Very rare.
 Snowy Owl, *Nyctea nyctea*. Very rare.
 Saw-whet Owl, *Nyctala acadica*. Rare.
 Purple Finch, *Carpodacus purpureus*. One record.
 Redpoll, *Acanthis linaria*. Very rare.
 Brown Creeper, *Certhia familiaris americana*. In extremely cold
 winters only, but then common.

REGULAR TRANSIENT VISITORS SOMETIMES SEEN IN
 WINTER, 14.

Loon, *Gavia imber*. Rare.
 American Merganser, *Merganser americana*. Common.
 Black Duck, *Anas obscura*. Common.
 Shoveller, *Spatula clypeata*. Rather rare.
 Pintail, *Dafla acuta*. Very common.
 American Golden-eye, *Clangula americana*. Rather rare.
 Canada Goose, *Branta canadensis*. Very Common.
 Wilson Snipe, *Gallinago delicata*. Generally stays all winter.
 Great Blue Heron, *Ardea herodias*. Rather rare.
 White-crowned Sparrow, *Zonotrichia leucophrys*. Common.
 Winter Wren, *Olbiorchilus hyemalis*. Rather rare.
 Red-breasted Nuthatch, *Sitta canadensis*. Very rare.
 Golden-crowned Kinglet, *Regulus satrapa*. Common.
 Ruby-crowned Kinglet, *Regulus calendula*. Rather rare.

ACCIDENTAL, 1.

Old-squaw, *Harelda hyemalis*.

INTRODUCED 2.

English Sparrow, *Passer domesticus*. Very common.
 Mongolian Pheasant, *Phasianus torquatus*. Rare.

WINTER BIRDS OF BRISTOL, CONN., AND VICINITY.

The birds here listed were recorded between January 1st and February 28th, 1902. While a list for a single winter should not be compared with one covering several winters, except as it indicates what species are pretty likely to be found during any winter, and therefore compares only with the species regularly found in that other locality, it is nevertheless interesting to notice the correspondences and differences which localities so far separated in longitude as well as latitude, from each other as these show. Other conditions, particularly the presence of the ocean and the intervening mountains, tend to emphasize the difference. It is possible here to give the actual number of records for each species for the two months covered. The notes are contributed by Mr. Frank Bruen.

Herring Gull, *Larus argentatus*. 9.
 Bob-white, *Colinus virginianus*. 50.
 Ruffed Grouse, *Bonasa umbellus*. 7.
 American Sparrow Hawk, *Falco sparverius*. 1.
 Pigeon Hawk, *Falco columbarius*. 3.
 Belted Kingfisher, *Ceryle alcyon*. 2.
 Hairy Woodpecker, *Dryobates villosus*. 1.
 Downy Woodpecker, *Dryobates pubescens medianus*. 18.
 Blue Jay, *Cyanocitta cristata*. 13.
 Northern Flicker, *Colaptes auratus luteus*. 1.
 American Crow, *Corvus americanus*. 220.
 Pine Siskin, *Spinus pinus*. (Possible error.)
 American Goldfinch, *Astraglinus tristis*. 150.
 Snoflake, *Passerina nivalis*. 300.
 Slate-colored Junco, *Junco hyemalis*. 30.
 Tree Sparrow, *Spizella monticola*. 150.
 Song Sparrow, *Melospiza melodia*. 25.
 Northern Shrike, *Lanius borealis*. 3.
 Winter Wren, *Olbiorchilus hyemalis*. 2.
 Brown Creeper, *Certhia familiaris americana*. 3.
 White-breasted Nuthatch, *Sitta carolinensis*. 36.
 Chickadee, *Parus atricapillus*. 53.
 Golden-crowned Kinglet, *Regulus satrapa*. 12.
 American Robin, *Merula migratoria*. 1.